Uterine Artery/Fibroid Embolization (UAE/UFE)

Preparation Instructions

You have been scheduled for a Uterine Artery/Fibroid Embolization (UAE/UFE).

This patient package contains the following:

- Information and preparation instructions for the procedure
- A map with directions to the Center
- Forms to fill out **before** your arrival:

Information/Details about your Condition

Patient Consultation Information (4 pages)

Discharge Instructions

Please bring these forms with you to the procedure.

- Expect to be at the office 3-4 hours.
- Arrange for someone to drive you home after the procedure.

Premier Radiology Briarville

1210 Briarville Road • Suite 602F • Madison, TN 37115

Hours: 7:30 AM - 5:00 PM

Office: 615-986-6411 Fax: 615-234-1506





Uterine Fibroid Embolization (UFE) Overview

In a UFE procedure, physicians use real time X-ray guidance to deliver small particles to the uterus and fibroids. The small particles are injected through a small tube called a catheter. These block the arteries that provide blood flow, causing the fibroids to shrink. Almost 90% of women with fibroids experience relief of their symptoms.

HOW SHOULD I PREPARE FOR THE PROCEDURE?

- Arrange for someone to drive you home after the procedure. You will be told to avoid driving for 24 hours as the narcotics that you receive will have to wear off.
- Discontinue any of the following medications for 3 days prior to your procedure: Plavix, Pletal, Ticlid, Trental, Coumadin, Warfarin, Brilinta, Xarelto or Pradaxa. Discontinue Effient for 7 days prior to your procedure. Patients who have had a drug eluting cardiac stent within the past year should NOT stop Plavix or Effient.
- Discontinue any of the following medicines 24 Hours prior to your procedure: Lovenox, Fragmin, Normiflo, Orgaran, Innohep, Arixtra, Eliquis, or Iprivask.
- All other medications are okay to take the day of the procedure. Take all your medications with small sips of water, unless told otherwise. It is very important to take blood pressure medications the morning of your examination. You will be told to stop taking blood thinners. This is important to prevent bleeding problems after your exam. Please bring all your medications with you to your appointment (not just a list). **DO NOT HOLD heart or blood pressure medications.**
- If you are on insulin, take ½ of your dose the morning of the procedure. Do not take any oral diabetic medication the morning of the procedure.
- Do not eat solid foods for **6 hours** prior to your procedure. You may drink clear liquids until **2 hours** before the procedure (no milk or creamer, no juice with pulp).
- If you have had imaging done prior to your procedure, please bring the disc and/or report with you.
- On the morning of your procedure please come to Briarville Premier Radiology office **30 minutes** before your appointment time. We open at 7:30 AM. (see attached map for directions).
- Inform the nurse and the radiologist of any allergies you have.
- Please fill out and bring the documents and questionnaire about your current health status, including current problems, allergies, medications, surgeries, etc.
- Nursing/technical staff will review your information that was completed and ask other questions. An IV will be started for IV sedation.
- The Interventional Radiologist will examine you and discuss the scheduled procedure. After questions are answered, we will have you sign a consent form. If you have any questions about the procedure before you arrive, you may call our office at **615-986-6411**. You will receive a reminder call the evening before the examination.





What Will Happen the Day of the Procedure

HOW IS THE PROCEDURE PERFORMED?

When you arrive at our office, you will check in at the reception desk. Paperwork will be verified and you will be escorted to the staging area. Our staff will ask several other questions. The Interventional Radiologist will explain the scheduled procedure, answer questions and perform a consultation. If the procedure will be performed the same day, you will receive some oral sedation and an intravenous (IV) line will placed in the arm. This allows us to give you fluids, antibiotics and/or medications as needed during your procedure. Once in the procedure room, you will be placed on your back and your groin area will be shaved and cleaned with a sterile solution which helps prevent infection. As you are sedated, monitoring devices will be placed on your chest and arms to evaluate your vital signs. Nursing staff will place a Foley catheter into your bladder. This is necessary in order to see the uterus during the procedure. Sedation will be provided to make sure that you are very comfortable and sleepy. A local anesthetic will be injected to numb the skin over the access site and a thin catheter (tube) will then be placed into the artery through your groin and directed with X-ray guidance into the uterine arteries. As contrast is injected, you may feel a sensation of warmth in the groin area and legs. This feeling will pass quickly. After injections determine proper catheter location, the arteries to the uterus will be blocked with microscopic particles on both the right and the left.

HOW LONG WILL THE PROCEDURE AND RECOVERY TAKE?

The uterine artery embolization procedure takes approximately 1 1/2 hours, sometimes longer if the anatomy is more complex. After the procedure is completed, the catheter will be removed and the Interventional Radiologist will seal the entry site with a closure device or stitch. Direct manual pressure is also applied for 5 to 10 minutes. This will help to prevent bruising and bleeding and allow the access site to heal. Typically you will be on bed rest for one hour after the procedure with your leg straight. Your pulse, blood pressure and entry site will be frequently evaluated during this period. You will be able to eat and drink. Some patients begin to notice some cramping in the early postoperative period. During recovery, your exam will be discussed with you and your family and any questions answered. The most difficult portion of the recovery is the night after the procedure. You will experience rather significant cramping like a very difficult menstrual period. Medications will be provided to help you through this portion of recovery.

WHEN WILL I KNOW THE RESULTS?

During the recovery period, the Interventional Radiologist will prepare some key images for you to take home at discharge. The exam and the images will be reviewed with you and your family. Prescriptions will be necessary, typically anti-inflammatory medications, antibiotics, steroids and pain medications. These prescriptions will be given to you at this time. Follow-up will be discussed as ultrasound/MRI exams are needed at 1, 3, 6 and 9 months. Reports will be generated for your referral doctor and primary care doctor. They will receive a copy of the consultation as well as the examination and treatment reports.

OUTPATIENT DISCHARGE INSTRUCTIONS

- Drink plenty of clear (nonalcoholic) liquids such as water, juice or soda once you have returned home at least 8 oz. every hour. You may begin your usual diet when you leave the Center.
- Resume your usual diet and any medications you routinely take (unless instructed not to do so by your doctor).
- For the next 12 hours, attempt to keep the leg in which the catheter was placed as straight as possible.
- No driving, operating heavy machinery, or making any legal decisions for 24 hours following the procedure.
- For several days after the procedure, avoid strenuous activities including vigorous exercise, housework, yard work, lifting objects heavier than 10 pounds or excessive bending at the waist.
- Do not take a tub bath until the groin site looks closed. This may be several days. You may shower.
- Please expect the Interventional Radiologist to call you the evening of the procedure to check on your status. A second follow-up call from the office will occur in several days following the procedure.
- It is very important that you take the provided medications at the time of discharge. These medications will prevent infection, reduce swelling of the uterus and help manage the discomfort that you will experience for the next several days.





Post-Uterine Fibroid Embolization Instructions

Activity:

- 1. For the next 12 hours, attempt to keep the leg in which the catheter was placed as straight as possible. This will help minimize your risk of bleeding (hematoma).
- 2. Go home and rest quietly for the remainder of the day. Do not drive, operate heavy machinery, or make any legal decisions for 24 hours.
- 3. The day after the procedure, avoid strenuous activities including vigorous exercise, lifting objects heavier than 10 pounds or excessive bending at the waist. This will help minimize your risk of bleeding (hematoma).

Diet/Medications:

- 1. You may resume your normal diet after the procedure.
- 2. Drink extra nonalcoholic fluids today (6-8 glasses). This will encourage the elimination of the contrast (dye).
- 3. You may restart all your medications (unless instructed not to do so by your doctor).
- 4. It is very important that you take the provided medications at the time of discharge. These medications will prevent infection, reduce swelling of the uterus and help manage the discomfort that you will experience for the next several days.
- 5. **Prescriptions:** Complete all as directed: Levaquin (antibiotic), Medrol dosepak (steroid). **Take as needed:** Toradol (non-narcotic pain medication), Lortab (for break-through pain), Stool softener, if taking Lortab for pain.

Wound Care:

- 1. Your procedure site should be kept clean and dry. Do not bathe in a tub for 48 hours. Showers are acceptable.
- 2. The dressing on your groin may be removed in the morning. Remove dressing while in the shower, replace with a band-aid for about 5 days.
- 3. Change the band-aid when it becomes soiled or wet.

Common Side Effects:

You should expect some discomfort for several days after your procedure. Take your usual pain medications prescribed by your Interventional Radiologist as directed.

Contact our office at 615-986-6411 for the following symptoms:

- 1. Bleeding or swelling at the entry site. If you notice swelling in your groin, apply direct pressure with your fingertips and call us immediately at **615-986-6411**. This may be a sign that bleeding has started from the artery or vein.
- 2. If you develop a fever above 101°, or any redness or drainage at the catheter site, or from the uterus, an infection could be developing. Seek medical attention immediately.
- 3. Any change in color or temperature of the leg. (A small bruise may be a normal occurrence.) If the leg in which the catheter was placed becomes cool, cold, numb or painful, call us immediately at **615-986-6411**, or the on-call service at **866-236-3572**. An Interventional Radiologist will be notified and will respond to your problem.

General Information:

- 1. You may have had a vascular closure device placed at the catheter site at the end of your procedure. Please refer to the information provided for specific details.
- 2. Call your referring doctor for a follow-up appointment. Results of your procedure will be sent directly to your referring doctor.
- 3. For any questions regarding your examination or problems you notice after leaving, please call Premier Radiology Briarville at 615-986-6411.


Thank you for allowing us to participate in your healthcare needs.

These discharge instructions have been explained to the patient. The patient or the person responsible for the patient fully understands these instructions.

Patient's Signature:	_ Date:	 Time:	
Signature of person giving instructions:			







FROM I-65 SOUTH FROM HENDERSONVILLE: On I-65 South take exit 92 Old Hickory Boulevard toward Madison. Turn left at the traffic light at the top of the ramp. Drive .4 miles to the traffic light at South Graycroft Avenue. Turn right onto South Graycroft Avenue. Continue straight for 1.1 miles to the intersection of South Graycroft Avenue and W. Due West Avenue. Turn right at the traffic light onto W. Due West Avenue. In 325 feet, turn left onto Briarville Road. (This is between the DCi building and the 607 Due West Medical Building). Turn right at the 4th driveway into the Briarville Medical Complex. Premier Radiology is in Building F with a blue awning over the door.

Premier Radiology Briarville: 1210 Briarville Road • Suite 602F • Madison, TN 37115 Hours: 7:30 AM - 5:00 PM • Office: 615-986-6411 • Fax: 615-234-1506

Information/Details about your Condition

Uterine Artery/Fibroid Embolization (UAE/UFE)

Menarche:	LMP:	Last I	PAP Smear:
Cycle Length:	Heavy Flo	w: Yes No	Clots with menses: Yes No
Number of Tampons:	Number of P	ads:	
Breast Tenderness: Yes No	Urinary Frequency:	Yes No Flui	d retention: Yes No
Abdominal bloating: Yes No	Pain with Intercourse:	☐ Yes ☐ No F	Fatigue: ☐ Yes ☐ No
Number of Pregnancies:	Number of Live Births:	Numbe	r of miscarriages/abortions:
Contraception: Pr	regnancy test today:	Negative	tive (Cancel procedure if positive)
Additional Gynecological history (la	paroscopy/hysteroscop	y/biopsy/neoplasr	m/fibroid surgery):
When were you first diagnosed with	n fibroids?		
Have you had recent imaging studi	es of the uterus/pelvis?	Yes No. If	Yes, when and where?
-			
Describe your symptoms (pain, full	ness, urinary frequency	etc):	
			· · · · · · · · · · · · · · · · · · ·
Describe how it feels (sharp, knife-	like, cramping, urgency,	burning, dull, ach	ing):
Rate your pain at its worst: 1 2 3	4 5 6 7 8 9 10 (circle	e)	
What do you do to help the pain? _		_ Do you take pai	n meds?
Do you have a history of anemia? [☐ Yes ☐ No. If yes, ha	s this been treate	d with medications or transfusions?





Patient Consultation Information

Date:			
Name: Da	ite of Birth:	_Age:	Sex
Referring MD:	Primary C	are MD:	
Phone contact information: Cell:		Home):
Describe why you are here today or what i	s your current prob	lem?	
Tell us more about your current problem: _			
ALLERGIES (please list medicine allergies): La	tex allergy? \(\text{YE}	S 🗆 NO	Nickel allergy? ☐ YES ☐ NO
Have you, or has anyone in your family, ev	er had a reaction to	o sedatior	n or anesthesia? ☐ YES ☐ NO
If yes, describe:			
CURRENT MEDICATIONS (BRING ALL MEDI	ICATIONS and list curre	ent medicat	ions and dosages below):
Date of last flu vaccine:			
SOCIAL HISTORY (things about you that will h	nelp us understand you	ır medical p	roblem): PLEASE CHECK
Smoking: ☐YES ☐ NO If yes, please d	escribe how long/h	ow much:	
If you have quit	smoking, when: _		
Alcohol: YES NO If you have quit	, when:		
Recreational drugs: YES NO Ir	n the past:		
Caffeine cups per day: Please give amoun	t:		
Marital status: (check one) ☐ Single ☐	Married 🗌 Widow	ved. Do	you live alone? ☐ YES ☐ NO
Occupation:			

Premier Radiology BRIARVILLE



PRIOR IMAGING: Have you na	id recent imaging exams (ditrasound, CT, MRT)?
Please list type and location: _	

PAST MEDICAL HISTORY (Medical care or problems):

Heart attack or rhythm problems	YES	NO
Pacemaker	YES	NO
Diabetes - treated with pills	YES	NO
Diabetes - treated with insulin	YES	NO
High blood pressure	YES	NO
Bleeding disorders	YES	NO
Stroke	YES	NO
Stomach or colon problems	YES	NO
Lung disease (cancer or emphysema)	YES	NO
Kidney disease	YES	NO
Mental/emotional disorder	YES	NO
Liver/jaundice/hepatitis	YES	NO
Vascular problems	YES	NO
Cancer	YES	NO
Musculoskeletal	YES	NO
Endocrine/Thyroid	YES	NO
Sleep Apnea	YES	NO
Other:		

REVIEW OF SYSTEMS (Past or current medical issues): Please circle

General:	Weight loss or gain	Fever	Chills
Neurological:	Headache	Confusion	Speech changes
Psychological:	Anxiety	Depression	
Skin:	Rash	Edema (swelling)	Bruising
Ears, Nose, Throat:	Hearing loss	Visual changes	Visual loss
Heart & Vessels:	Chest pain	Irregular heart beat	Leg edema (swelling)
Lungs & Breathing:	Problems at rest	Problem with exercise	Problem with lying flat
Stomach & Bowels:	Vomiting or nausea	Diarrhea or constipation	Blood in stool
Kidneys & Bladder:	Frequent urination	Painful urination	Bloody urine
Muscles & Bones:	Leg pain with movement-R or L	Leg pain with rest-R or L	Arthritis
Other:			
Other:			

FAMILY HISTORY: (Please list all known major family medical history).

Problem	Father	Mother	Brother or Sister	Describe
Heart Disease				
Stroke				
Abdominal Aneurysm				
High Blood Pressure				
Cancer (location under other)				
Other:				
Other:				
Deceased or Alive? Age?				

PAST SURGICAL HISTORY: (Please list prior surgeries with approximate dates).

			Dates
Heart surgery	YES	NO	
Back surgery	YES	NO	
Knee or hip surgery	YES	NO	
Aortic surgery (aneurysm treatment)	YES	NO	
Leg bypass surgery for artery disease	YES	NO	
Carotid artery surgery (neck artery)	YES	NO	
Aneurysm treatment with coils inside of head	YES	NO	
Gallbladder or colon	YES	NO	
Uterus and/or ovaries	YES	NO	
Other:			
Other:			
Other:			

How did you learn about Premier Radiology Briarville?

T:	P:	R:	BP(R):	BP(I	L):
Oxygen Sat%:		on room a	nir. Oxygen Sat%	on:	LPN
Weight:		Height:	BMI:		
Labs: Cr/GFR:		/ PT/I	INR:/	Blood Glucos	se:
Mental Status:	□Ale	rt	Confused Defic	cits:	learing
Airway:	□No	rmal 🗌 Limited ne	eck movement	Dentures ☐ Caro	otid Bruit: R L
Lungs:	☐ Cle	ear 🗌 Rales 🗀	Rhonchi Whe	ezes	
Heart:	Re	gular 🗌 Murmur			
Abdomen:	Bowe	I Sounds ☐ Yes ☐	No Masses □	Yes □ No Bruit	∷ ∐Yes
Leg strength:	□No	rmal 🗌 Weak			· · · · · · · · · · · · · · · · · · ·
Arm strength:	□No	rmal 🗌 Weak			
Patellar Reflex	es:				
Pain with strai	ght leg	raises?: Yes	No		
Pre-procedure	Pulses	s:			
Right		Femoral	PT	DP	Radial
Value: +3 to do	ppler			21	
		Femoral Femoral	PT PT	DP DP	Radial Radial
Value: +3 to do Left Value: +3 to do	ppler	Femoral	PT	DP	Radial
Value: +3 to do Left Value: +3 to do Women: Date of	ppler of last m	Femoral nenstrual period:	PT	DP Pregnancy possib	Radial
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati	ppler of last m	Femoral nenstrual period:	PT platelet agents: Ye	DP Pregnancy possib	Radial ility? Yes No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available	ppler of last maion of before train	Femoral nenstrual period: lood thinners or anti	PT ———————————————————————————————————	DP Pregnancy possib	Radial liity? ☐ Yes ☐ No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available Completed by:	ppler of last maion of be	Femoral nenstrual period: lood thinners or anti- nsportation: \[\sum Yes \]	PT platelet agents: ☐ Ye No	Pregnancy possib	Radial ility? Yes No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available Completed by:	ppler of last maion of be	Femoral nenstrual period: lood thinners or anti- nsportation: \[\sum Yes \]	PT ———————————————————————————————————	Pregnancy possib	Radial ility? Yes No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available Completed by:	ppler of last maion of be	Femoral nenstrual period: lood thinners or anti- nsportation: \[\sum Yes \]	PT platelet agents: ☐ Ye No	Pregnancy possib	Radial ility? Yes No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available Completed by:	ppler of last maion of be	Femoral nenstrual period: lood thinners or anti- nsportation: \[\sum Yes \]	PT platelet agents: ☐ Ye No	Pregnancy possib	Radial ility? Yes No
Value: +3 to do Left Value: +3 to do Women: Date of Confirm cessati Driver available Completed by:	ppler of last man	Femoral nenstrual period: lood thinners or anti	PT platelet agents: ☐ Ye No	Pregnancy possib	Radial ility? Yes No